

zpoet	Pr1	Pr2	Pr3	Pr4	Pr5	Pr6	Pr7	Σ	znmka

v poliach R a C

1. [13 bodov] Dan je matica $A = \begin{pmatrix} 1 & 2 & 1 \\ 1 & 1 & -1 \\ -2 & 3 & 4 \end{pmatrix}$ a $\mathbf{b} = \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}$. Urte
- a) [1] stopu matice A , b) [3] minimlny polynm $m_{A,\mathbf{b}}(\lambda)$ (vektora $\mathbf{b}$ vzhadom na maticu A),
c) [3] vlastn sla matice A , d) [4] Jordanovu maticu J a maticu P , pre ktor $A = PJP^{-1}$,
e) [2] maticu $f(J)$, pre $f(\lambda) = (\lambda - 2)^9 + \lambda^2 - 1$.
2. [8] Njdite rovnicu priamky $y(x) = kx + q$, pre ktor je set
 $\sum_{i=-2}^2 |y(x_i) - y_i|^2$ najmen. Hodnoty y_i s v tabuke $\rightarrow$
- | | | | | | |
|-------|----|----|---|---|---|
| x_i | -2 | -1 | 0 | 1 | 2 |
| y_i | -1 | 0 | 2 | 3 | 4 |
3. [10] Nech $A = \begin{pmatrix} 2 & -2 & -1 & 0 & 2 \\ 1 & -1 & 1 & 0 & 2 \\ 0 & 0 & 6 & 3 & 1 \\ 1 & -1 & 1 & 1 & 1 \end{pmatrix} \in R^{4 \times 5}$. $L_r(A) = \text{span}\{A_{1*}, A_{2*}, A_{3*}, A_{4*}\}$ je riadkov,
 $L_s(A) = \text{span}\{A_{*1}, A_{*2}, A_{*3}, A_{*4}, A_{*5}\}$ stpcov a $N(A) = \{\mathbf{x} \in R^{5 \times 1} : A\mathbf{x} = 0\}$ nulov priestor matice A .
Urte bzy a dimenzie priestorov $L_r(A)$, $L_s(A)$ a $N(A)$.
4. Nech L , M s linerne priestory nad poom R , $\mathcal{B} = \{\mathbf{b}_1, \mathbf{b}_2, \mathbf{b}_3, \mathbf{b}_4, \mathbf{b}_5\}$ je bza priestoru L , $\mathcal{D} = \{\mathbf{d}_1, \mathbf{d}_2, \mathbf{d}_3\}$ je bza priestoru M . Napte
- a) (1 bod) $\dim M$ b) (1 bod) sradnice $[2\mathbf{b}_1 - \mathbf{b}_3 + 3\mathbf{b}_4]_{\mathcal{B}}$
c) [3] maticu $T_{\mathcal{B}\mathcal{D}}$ linerneho opertora T urenho vazmi
 $T\mathbf{b}_1 = \mathbf{d}_2$, $T\mathbf{b}_2 = \mathbf{d}_1 + \mathbf{d}_2 + \mathbf{d}_3$, $T\mathbf{b}_3 = 2\mathbf{d}_1 + \mathbf{d}_2 + 2\mathbf{d}_3$, $T\mathbf{b}_4 = 0$, $T\mathbf{b}_5 = -\mathbf{d}_1 + 2\mathbf{d}_2 - \mathbf{d}_3$.
d) [3] $\dim \ker T$ a $\dim \text{ran } T$

V konenich poliach

5. [6 bodov] Pomocou Euklidovho algoritmu zistite, či má polynóm $f(x) = x^4 + x^3 + x + 1 \in P(Z_3)$ koreň násobnosti aspoň 2.
6. [10] V poli F riete sstavu linernych rovnc
- | | | | |
|--------------|-----------------------|--------------|------------------------|
| a) $F = Z_2$ | $x_1 + x_3 + x_4 = 1$ | b) $F = Z_3$ | $x_1 + 2x_3 + x_4 = 1$ |
| | $x_2 + x_3 = 0$ | | $2x_2 + x_3 = 2$ |
| | $x_1 + x_2 + x_4 = 1$ | | $x_1 + x_2 + 2x_4 = 1$ |
7. a) [2] Napte koko prvkov m pole $F = P(Z_2)/(x^3 + x + 1)$.
b) [3] Vypotajte $(x+1)(x^2+x) \pmod{x^3+x+1}$.
c* [3] Dokte tvrdenie: Ak je $f(x) \in P(F)$ ireducibln polynm, tak m kad prvok $0 \neq h \in P(F)/f(x)$ inverzn (t.j. $\exists g \in P(F)$, pre ktor $g(x)h(x) = 1 \pmod{f(x)}$).

Riešenie.

1a. $\text{trace}(A) = 6$,

b) $A = \begin{pmatrix} 1 & 2 & 1 \\ 1 & 1 & -1 \\ -2 & 3 & 4 \end{pmatrix}$, $Ab = \begin{pmatrix} 3 \\ 2 \\ 1 \end{pmatrix}$, $A^2b = A(Ab) = \begin{pmatrix} 8 \\ 4 \\ 4 \end{pmatrix}$, $A^3b = A(A^2b) = \begin{pmatrix} 20 \\ 8 \\ 12 \end{pmatrix}$,
 $\begin{pmatrix} 1 & 3 & 8 & 20 \\ 1 & 2 & 4 & 8 \\ 0 & 1 & 4 & 12 \end{pmatrix} \xrightarrow[R_2 - R_1]{R_1 \leftrightarrow R_2} \sim \begin{pmatrix} 1 & 2 & 4 & 8 \\ 0 & 1 & 4 & 12 \\ 0 & 1 & 4 & 12 \end{pmatrix}$, $a_3 = 0, a_2 = 1$, t.j. $m_{A,b}(\lambda) = \lambda^2 - 4\lambda + 4 = (\lambda - 2)^2$
 $a_0 = 4$

c) $\lambda_{1,2} = 2, 2 + 2 + \lambda_3 = \text{trace}(A) = 6 \implies \lambda_{1,2,3} = 2$

d) Z rovnice $(A - 2I)P_{*3} = 0$ dostaneme $P_{*3} = (1 \ 0 \ 1)^\top$, potom rieime sstavu $(A - 2I)P_{*2} = P_{*3}$ a $(A - 2I)P_{*1} = P_{*2}$:

$$\left(\begin{array}{ccc|cc} -1 & 2 & 1 & 1 & 1 \\ 1 & -1 & -1 & 0 & 1 \\ -2 & 3 & 2 & 1 & 0 \end{array} \right) \xrightarrow[R_3 + 2R_2]{R_1 + R_2} \sim \left(\begin{array}{ccc|cc} 0 & 1 & 0 & 1 & 2 \\ 1 & -1 & -1 & 0 & 1 \\ 0 & 1 & 0 & 1 & 2 \end{array} \right), \quad P = \begin{pmatrix} 1 & 1 & 1 \\ 2 & 1 & 0 \\ -2 & 0 & 1 \end{pmatrix}, \quad J = \begin{pmatrix} 2 & 0 & 0 \\ 1 & 2 & 0 \\ 0 & 1 & 2 \end{pmatrix}$$

e) $f(2) = 3; f'(\lambda) = 9(\lambda - 2)^8 + 2\lambda, f'(2) = 4; f''(\lambda) = 72(\lambda - 2)^7 + 2, \frac{f''(2)}{2!} = 1 \implies f(J) = \begin{pmatrix} 3 & 0 & 0 \\ 4 & 3 & 0 \\ 1 & 4 & 3 \end{pmatrix}$

2. $A = \begin{pmatrix} -2 & 1 \\ -1 & 1 \\ 0 & 1 \\ 1 & 1 \\ 2 & 1 \end{pmatrix}, \mathbf{b} = \begin{pmatrix} -1 \\ 0 \\ 2 \\ 3 \\ 4 \end{pmatrix}. A^\top A = \begin{pmatrix} 10 & 0 \\ 0 & 5 \end{pmatrix}, A^\top \mathbf{b} = \begin{pmatrix} 13 \\ 8 \end{pmatrix}.$

$$A^\top A \begin{pmatrix} k \\ q \end{pmatrix} = A^\top \mathbf{b} \implies k = 1, 3, q = 1, 6, \underline{y(x) = 1, 3x + 1, 6}.$$

3.

$$\begin{pmatrix} 2 & -2 & -1 & 0 & 2 \\ 1 & -1 & 1 & 0 & 2 \\ 0 & 0 & 6 & 3 & 1 \\ 1 & -1 & 1 & 1 & 1 \end{pmatrix} \sim \begin{pmatrix} 1 & -1 & 1 & 0 & 2 \\ 0 & 0 & 3 & 0 & 2 \\ 0 & 0 & 0 & 1 & -1 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix} = B$$

$$\mathcal{B}_r = \{B_{1*}, B_{2*}, B_{3*}\} = \{(1, -1, 1, 0, 2); (0, 0, 3, 0, 2); (0, 0, 0, 1, -1)\}, \dim L_r(A) = 3 = \dim L_s(A)$$

$$\mathcal{B}_s = \{A_{1*}, A_{3*}, A_{4*}\} \text{ (stĺpce matice } A, \text{ v ktorých je upravená matrica pivoty)}, \dim N(A) = 5 - 3 = 2,$$

$$\mathcal{B}_{N(A)} = \{(-4, 0, -2, 3, 3)^\top; (1, 1, 0, 0, 0)^\top\} \text{ (báza riešenia stavu } Ax = 0).$$

4. a) $\dim M = 3$, b) $[2\mathbf{b}_1 - \mathbf{b}_3 + 3\mathbf{b}_4]_{\mathcal{B}} = (2, 0, -1, 3, 0)^\top$, c) $T_{\mathcal{BD}} = \begin{pmatrix} 0 & 1 & 2 & 0 & -1 \\ 1 & 1 & 1 & 0 & 2 \\ 0 & 1 & 2 & 0 & -1 \end{pmatrix}$,

d) $\dim \ker T = 3, \dim \text{ran } T = 2$.

5. $f_1(x) = x^4 + x_3 + x + 1, f_2(x) = f'(x) = x^3 + 1$

$$(x^4 + x_3 + x + 1) : (x^3 + 1) = x + 1 \implies \gcd(f, f') = x^3 + 1, c = -1 \text{ je koreň polynómu } f' \implies \text{polynóm } f \text{ má koreň násobnosti } \geq 2.$$

6a. $P = \{(1, 0, 0, 0); (0, 0, 0, 1); (0, 1, 1, 0); (1, 1, 1, 1)\}$

6b. $P = \{(2, 2, 1, 0); (2, 0, 2, 1); (2, 1, 0, 2)\}$

7a. $2^3 = 8$

7b. $(x + 1)(x^2 + x) = x(x + 1)^2 = x^3 + x = (x^3 + x + 1) + 1 = 1 \pmod{(x^3 + x + 1)}$ (v $P(Z_2)$)

7c* $h \not\equiv 0 \pmod{f(X)}, \text{stup } h < \text{stup } f \implies \gcd(h, f) = 1$.

Teda $\exists a(x), b(x)$, pre ktor $1 = a(x)h(x) + b(x)f(x) = a(x)h(x) \pmod{f(x)}$, t.j. $g(x) = a(x)$.